Supplementary data for the article:

Waisi, H.; Janković, B.; Nikolić, B.; Dragičević, V.; Panić, I.; Tosti, T.; Trifković, J.
Influence of Various Concentrations of 24-Epibrassinolide on the Kinetic Parameters
during Isothermal Dehydration of Two Maize Hybrids. *South African Journal of Botany* **2018**, *119*, 69–79. <u>https://doi.org/10.1016/j.sajb.2018.08.006</u>

South African Journal of Botany

Influence of various concentrations of 24-epibrassinolide on the kinetic

parameters during isothermal dehydration of two maize hybrids

Authors: Hadi Waisi^{a,b}*, Bojan Janković, Bogdan Nikolić, Vesna Dragičević, Ivan Panić, Tomislav Tosti, Jelena Trifković

**Corresponding Author affiliation*:

^aFaculty for Ecology and Environmental Protection, University Union-Nikola Tesla, Cara Dušana 62-64, 11000, Belgrade, Serbia ^bInstitute of Nuclear Sciences "Vinča", University of Belgrade, Department of Physical Chemistry, Mike Petrovica Alasa 12-14, 11351 Vinca, Belgrade, Serbia


Content of Supplementary Material:

Fig. S.1. Isothermal mass loss experiments ($\Delta T = 60 - 130$ °C) for dehydration process of the control samples for the ZP434 and ZP704 hybrid maize systems, including all seedling parts: radicle ((a), (d)), plumule ((b), (e)), and RoS ((c), (f)), respectively.

Fig. S.2. Isothermal mass loss experiments ($\Delta T = 60 - 130$ °C) for dehydration process of ZP434 (red designations) and ZP704 (blue designations) hybrids, including all seedling parts, treated with various concentrations of 24-EBL (5.20×10^{-9} M, 5.20×10^{-12} M, and 5.20×10^{-15} M, respectively).

Fig. S.3. Reduced time plots for selected reaction models, considering dehydration process for control samples of both maize hybrids.

Fig. S.4. Reduced time plots for selected reaction models, including all seedling parts (radicle, plumule and RoS) treated with various concentrations of 24-EBL (5.20×10^{-9} M, 5.20×10^{-12} M and 5.20×10^{-15} M).







b


e

h





f


i



























n


0

















Fig. S.2.



d










Fig. S.3.






i























g






h







0

r
































р








Fig. S.4.