

Supplementary material for the article:

Samelak, I.; Balaban, M.; Vidović, N.; Koljančić, N.; Antić, M.; Šolević-Knudsen, T.; Jovančičević, B. Application of Alkane Biological Markers in the Assessment of the Origin of Oil Pollutants in the Soil and Recent River Sediments (River Vrbas, Bosnia and Herzegovina). *Journal of the Serbian Chemical Society* **2018**, *83* (10), 1167–1175.  
<https://doi.org/10.2298/JSC180501061S>



*J. Serb. Chem. Soc.* 83 (10) S322–S325 (2018)

SUPPLEMENTARY MATERIAL TO  
**Application of alkane biological markers in the assessment of the  
origin of oil pollutants in the soil and recent river sediments  
(River Vrbas, Bosnia and Herzegovina)**

IVAN SAMELAK<sup>1</sup>, MILICA BALABAN<sup>1\*</sup>, NADA VIDOVIĆ<sup>1</sup>, NEMANJA KOLJANČIĆ<sup>1</sup>,  
MALIŠA ANTIC<sup>2</sup>, TATJANA ŠOLEVIĆ-KNUDSEN<sup>3</sup> and BRANIMIR JOVANČIĆEVIĆ<sup>4</sup>

<sup>1</sup>University of Banja Luka, Faculty of Natural Sciences and Mathematics, Mladena Stojanovića 2, 78000 Banja Luka, Bosnia and Herzegovina, <sup>2</sup>University in Belgrade, Faculty of Agriculture, Nemanjina 6, 11080 Zemun, Serbia, <sup>3</sup>University of Belgrade, Center of Chemistry, Institute of Chemistry, Technology and Metallurgy, Njegoševa 12, 11001 Belgrade, Serbia and <sup>4</sup>University of Belgrade, Faculty of Chemistry, Studentski trg 12–16, 11001 Belgrade, Serbia

*J. Serb. Chem. Soc.* 83 (10) (2018) 1167–1175

SAMPLES

The studied samples of fresh river sediments and bank sediments (soils) belong to the Vrbas River. The samples were taken near the sewerage drainage in the central zone of the city of Banja Luka. The watershed of the Vrbas River and the locations of the sampling sites of the four samples of fresh river sediment (RS<sub>1</sub>–RS<sub>4</sub>) and the four samples of coastal sediments, *i.e.*, soils (S<sub>1</sub>–S<sub>4</sub>) are shown in Fig. S-1. The fresh river sediments were taken at the boundary of water and the bank, while the soils were taken at a few meters distance from the River. All samples were taken at a depth of 10 cm.

\* Corresponding author. E-mail: milica.balaban@pmf.unibl.org



Fig. S-1. Flow of the Vrbas River with the sampling locations of the river sediments, RS<sub>1</sub>–RS<sub>4</sub>, and the riverbank sediments (soils), S<sub>1</sub>–S<sub>4</sub>.

TABLE S-I. Identification of representative sterane and terpane peaks in fragmentograms  $m/z$  217 and  $m/z$  191 shown in Fig. S-1b and c

Peak	Compounds
1	C <sub>27</sub> 13 $\beta$ (H)17 $\alpha$ (H)20(S)-diasterane
2	C <sub>28</sub> 13 $\alpha$ (H)17 $\beta$ (H)20(R)-diasterane + C <sub>27</sub> 14 $\beta$ (H)17 $\beta$ (H)20(S)-sterane
3	C <sub>27</sub> 14 $\alpha$ (H)17 $\alpha$ (H)20(R)-sterane
4	C <sub>28</sub> 14 $\beta$ (H)17 $\beta$ (H)20(S)-sterane
5	C <sub>28</sub> 14 $\alpha$ (H)17 $\alpha$ (H)20(R)-sterane
6	C <sub>29</sub> 14 $\beta$ (H)17 $\beta$ (H)20(S)-sterane
7	C <sub>29</sub> 14 $\alpha$ (H)17 $\alpha$ (H)20(R)-sterane
8	C <sub>29</sub> 17 $\alpha$ (H)21 $\beta$ (H)-hopane
9	C <sub>30</sub> 17 $\alpha$ (H)21 $\beta$ (H)-hopane
10	C <sub>31</sub> 17 $\alpha$ (H)21 $\beta$ (H)-hopane (22S and 22R)
11	C <sub>32</sub> 17 $\alpha$ (H)21 $\beta$ (H)-hopane (22S and 22R)
12	C <sub>33</sub> 17 $\alpha$ (H)21 $\beta$ (H)-hopane (22S and 22R)
13	C <sub>34</sub> 17 $\alpha$ (H)21 $\beta$ (H)-hopane (22S and 22R)
14	C <sub>35</sub> 17 $\alpha$ (H)21 $\beta$ (H)-hopane (22S and 22R)




Fig. S-2. Fragmentograms of *n*-alkanes ( $m/z$  71), steranes ( $m/z$  217) and terpanes ( $m/z$  191) of the RS<sub>1</sub> and S<sub>1</sub> samples, which are typical for the extracts of the investigated fresh river sediments and soils.




Fig. S-3. Fragmentograms of *n*-alkanes ( $m/z$  71), steranes ( $m/z$  217) and terpanes ( $m/z$  191) of the river sediments samples, RS<sub>1</sub>-RS<sub>4</sub> and bank sediments (soils) samples, S<sub>1</sub>-S<sub>4</sub>.