

Supplementary data for the article:

Pantelić, M. M.; Dabić Zagorac, D. Č.; Davidović, S. M.; Todić, S. R.; Bešlić, Z. S.; Gašić, U. M.; Tešić, Ž. L.; Natić, M. M. Identification and Quantification of Phenolic Compounds in Berry Skin, Pulp, and Seeds in 13 Grapevine Varieties Grown in Serbia. *Food Chemistry* **2016**, *211*, 243–252. <https://doi.org/10.1016/j.foodchem.2016.05.051>

**Identification and quantification of phenolic compounds in berry skin, pulp, and seeds
in 13 grapevine varieties grown in Serbia**

Milica M. Pantelić^a, Dragana Č. Dabić Zagorac^a, Sonja M. Davidović^b, Slavica R. Todić^b, Zoran
S. Bešlić^b, Uroš M. Gašić^c, Živoslav Lj. Tešić^c and Maja M. Natić^{c*}

^aInnovation Center, Faculty of Chemistry Ltd, University of Belgrade, 11158 Belgrade, Serbia

^bFaculty of Agriculture, University of Belgrade, Nemanjina 6, 11080 Belgrade, Serbia

^cFaculty of Chemistry, University of Belgrade, P. O. Box 51, 11158 Belgrade, Serbia

*Corresponding author:

Maja M. Natić

E-mail: mnatic@gmail.com

Address: Studentski trg 12-16, 11000 Belgrade, Serbia

Tel: +381113336730

Fax: +381112635357

Table S1. Intensities obtained from the full scan spectra for identified anthocyanins in **studied** grape skin samples.

	Cabernet Sauvignon	Merlot	Cabernet Franc	Sangiovese	Shiraz	Pinot noir	Prokupac
Delphinidin 3- <i>O</i> -hexoside isomer 1	53.33	90.07	67.19	32.46	6.50	112.17	6.64
Cyanidin 3- <i>O</i> -hexoside isomer 1	15.14	14.66	20.53	55.15	2.60	15.34	1.91
Petunidin 3- <i>O</i> -hexoside isomer 1	63.39	75.62	98.73	37.90	12.65	136.13	13.52
Malvidin 3,5- <i>O</i> -dihexoside	0.68	0.19	1.47	0.08	0.92	2.24	0.88
Peonidin 3- <i>O</i> -hexoside	95.84	70.29	164.69	33.98	60.86	78.55	30.31
Malvidin 3- <i>O</i> -hexoside	401.58	393.04	462.74	60.27	126.98	569.47	159.35
Petunidin 3- <i>O</i> -acetyl hexoside	0.30	0.36	0.34	–	0.30	–	0.08
Malvidin 3- <i>O</i> -acetyl hexoside isomer 1	1.73	1.91	2.18	–	0.99	–	0.13
Peonidin 3- <i>O</i> -acetyl hexoside	0.63	0.32	0.57	–	0.11	–	0.08
Delphinidin 3- <i>O-p</i> -coumaroyl hexoside	2.00	2.22	13.61	0.21	0.30	19.60	0.57
Malvidin 3- <i>O</i> -acetyl hexoside isomer 2	9.00	5.45	4.36	–	2.53	–	0.92
Delphinidin 3- <i>O</i> -hexoside isomer 2	24.79	12.50	57.05	14.39	0.26	16.12	3.38
Delphinidin 3- <i>O</i> -hexuronide	5.87	3.00	4.84	2.55	0.35	5.19	0.88
Malvidin 3- <i>O</i> -caffeoyl hexoside	2.26	1.77	3.97	–	–	13.10	0.78
Cyanidin 3- <i>O-p</i> -coumaroyl hexoside	0.80	0.65	8.52	0.92	0.13	3.16	0.21
Petunidin 3- <i>O-p</i> -coumaroyl hexoside	4.20	4.32	31.38	0.25	0.41	33.06	1.47
Peonidin 3- <i>O-p</i> -coumaroyl hexoside	18.88	12.84	97.55	0.40	0.94	29.62	6.42
Cyanidin 3- <i>O</i> - hexoside isomer 2	3.02	1.06	4.48	0.60	–	2.37	0.77
Malvidin 3- <i>O-p</i> -coumaroyl hexoside	101.59	84.48	239.93	0.83	3.31	270.49	44.42
Petunidin 3- <i>O</i> -hexoside isomer 2	7.37	1.65	25.02	0.13	0.14	2.48	2.40