

Supplementary data for article:

Casey, W. T.; Nikodinović-Runić, J.; Fonseca Garcia, P.; Guzik, M. W.; McGrath, J. W.; Quinn, J. P.; Cagney, G.; Auxiliadora Prieto, M.; O'Connor, K. E. The Effect of Polyphosphate Kinase Gene Deletion on Polyhydroxyalkanoate Accumulation and Carbon Metabolism in *Pseudomonas Putida* KT2440. *Environmental Microbiology Reports* 2013, 5 (5), 740–746.

<https://doi.org/10.1111/1758-2229.12076>

Supporting Table 3.

Bacterial strains used in this study

Strain	Relevant Characteristics	Source
<i>P. putida</i> KT2440	Wild-type strain, amp ^R	ATTC 47054
<i>P. putida</i> KT2440 Δppk	<i>P. putida</i> KT2440 ppk knockout mutant, amp ^R , gen ^R	<i>This study</i>
<i>P. putida</i> KT2440 Δppk/pJB861ppk	<i>P. putida</i> KT2440 ppk knockout mutant complement, expressing ppk from pJB861ppk, amp ^R , kan ^R , gen ^R	<i>This study</i>
<i>Escherichia coli</i> DH5α	<i>F</i> -, <i>U169</i> , <i>recA1</i> , <i>endA1</i> , <i>tonA</i> , <i>hdR17</i> , <i>gyrA96</i> , <i>relA1</i> , cloning host	Life Technologies
<i>Escherichia coli</i> XL10 Gold	<i>endA1</i> , <i>recA1</i> , <i>gyrA96</i> , <i>relA1</i> , <i>lac</i> , Tet ^R , cloning host	Stratagene