

Supplementary data for the article:

Uzelac, T. N.; Nikolić-Kokić, A. L.; Spasić, S. D.; Mačvanin, M. T.; Nikolić, M. R.; Mandić, L. M.; Jovanović, V. B. Opposite Clozapine and Ziprasidone Effects on the Reactivity of Plasma Albumin SH-Group Are the Consequence of Their Different Binding Properties Dependent on Protein Fatty Acids Content. *Chemico-Biological Interactions* **2019**, *311*. <https://doi.org/10.1016/j.cbi.2019.108787>

## Electronic Supplementary Information

### **Opposite clozapine and ziprasidone effects on the reactivity of plasma albumin SH-group are the consequence of their different binding properties dependent on protein fatty acids content**


Tamara N. Uzelac <sup>a</sup>, Aleksandra L. Nikolić-Kokić <sup>b</sup>, Snežana D. Spasić <sup>c</sup>, Mirjana T. Mačvanin <sup>a</sup>,  
Milan R. Nikolić <sup>a</sup>, Ljuba M. Mandić <sup>a</sup>, and Vesna B. Jovanović <sup>a\*</sup>

<sup>a</sup> *University of Belgrade - Faculty of Chemistry, Department of Biochemistry, and Center of Excellence for Molecular Food Sciences, Studentski trg 12-16, 11000 Belgrade, Serbia*


<sup>b</sup> *University of Belgrade, Institute for Biological Research "Siniša Stanković", Department of Physiology, Bulevar despota Stefana 142, 11000 Belgrade, Serbia*

<sup>c</sup> *University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Centre for Chemistry, Njegoševa 12, 11000 Belgrade, Serbia*


\*Corresponding author: Dr. Vesna B. Jovanović, University of Belgrade - Faculty of Chemistry, Studentski trg 12-16, 11000 Belgrade, Serbia; Phone number: +381 11 333 66 76; E-mail: vjovanovic@chem.bg.ac.rs (V. Jovanović).



**Fig. S1.** Quenching of (de)fattened human serum albumin (HSA; 1  $\mu\text{M}$ ) fluorescence by clozapine (CLZ) or ziprasidone (ZIP) (0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4 and 4.5  $\mu\text{M}$ ) in 0.1 M sodium phosphate buffer pH 7.4 at 37°C (excitation wavelength 295 nm); HSA/SA, HSA in complex with stearic acid.



**Fig. S2.** Stern-Volmer plots of (de)fatted human serum albumin (HSA) fluorescence quenched by clozapine (CLZ) or ziprasidone (ZIP) at 37°C and pH 7.4; HSA/SA, HSA in complex with stearic acid (molar ratios from 1:1 up to 1:4). Error bars represent the standard deviation (n=2).



**Fig. S3.** Fluorescence quenching based plots, from data for the determination of binding constants and the number of binding sites of (de)fatted human serum albumin (HSA) in complex with clozapine (CLZ) or ziprasidone (ZIP) at 37°C and pH 7.4; HSA/SA, HSA in complex with stearic acid (molar ratios from 1:1 up to 1:4). Error bars represent the standard deviation (n=2).