

Supplementary data for article:

Zianna, A.; Sumar Ristic, M.; Psomas, G.; Hatzidimitriou, A.; Coutouli-Argyropoulou, E.; Lalia-Kantouri, M. Cadmium(II) Complexes of 5-Nitro-Salicylaldehyde and α - Diimines: Synthesis, Structure and Interaction with Calf-Thymus DNA. *Journal of Coordination Chemistry* **2015**, 68 (24), 4444–4463.

<https://doi.org/10.1080/00958972.2015.1101075>

Cadmium(II) complexes of 5-nitro-salicylaldehyde and α -diimines: Synthesis, structure and interaction with calf-thymus DNA

Ariadni Zianna ^a, Maja Sumar Ristic ^{a, b}, George Psomas ^a, Antonios G. Hatzidimitriou ^a, Evdoxia Coutouli-Argyropoulou ^c, Maria Lalia-Kantouri ^{a *}

^a Department of General and Inorganic Chemistry, Faculty of Chemistry, Aristotle University of Thessaloniki, GR-54124 Thessaloniki, Greece

^b Faculty of Chemistry, University of Belgrade, Studenski trg 12-16, Belgrade, Serbia

^c Department of Organic Chemistry and Biochemistry, Faculty of Chemistry, Aristotle University of Thessaloniki, GR-54124 Thessaloniki, Greece

S1. Interaction with CT DNA

The binding constant, K_b , can be obtained by monitoring the changes in the absorbance at the corresponding λ_{\max} with increasing concentrations of CT DNA and it is given by the ratio of slope to the y intercept in plots $\frac{[\text{DNA}]}{(\varepsilon_A - \varepsilon_f)}$ versus [DNA], according to the Wolfe–Shimer equation

[S1]:

$$\frac{[\text{DNA}]}{(\varepsilon_A - \varepsilon_f)} = \frac{[\text{DNA}]}{(\varepsilon_b - \varepsilon_f)} + \frac{1}{K_b(\varepsilon_b - \varepsilon_f)} \quad (\text{eq. S1})$$

where [DNA] is the concentration of DNA in base pairs, $\varepsilon_A = A_{\text{obsd}}/[\text{compound}]$, ε_f = the extinction coefficient for the free compound and ε_b = the extinction coefficient for the compound in the fully bound form.

S2. Competitive studies with EB

The Stern–Volmer constant K_{SV} is used to evaluate the quenching efficiency for each compound according to the Stern–Volmer equation [S2]:

$$\frac{I_0}{I} = 1 + K_{SV}[Q] \quad (\text{eq. S2})$$

* Corresponding author. Tel./fax: +30 2310 997844. E-mail address: lalia@chem.auth.gr (M. Lalia-Kantouri).

where I_0 and I are the emission intensities in the absence and the presence of the quencher, respectively, $[Q]$ is the concentration of the quencher (i.e. complexes **1–5**); K_{SV} is obtained from the Stern–Volmer plots by the slope of the diagram $\frac{I_0}{I}$ vs $[Q]$.

References

- [S1] A. Wolfe, G. Shimer, T. Meehan, *Biochemistry*, **26**, 6392 (1987).
[S2] J.R. Lakowicz, *Principles of Fluorescence Spectroscopy*, 3rd Edn., Plenum Press, New York (2006).




Fig. S1. (A)–(E) Plot of $\frac{[DNA]}{(\epsilon_A - \epsilon_f)}$ vs $[DNA]$ for complexes **1–5**, respectively.




Fig. S2. (A)–(E) Stern–Volmer quenching plot of EB bound to CT DNA for complexes **1–5**, respectively.