




1st FoodEnTwin Workshop “Food and Environmental -Omics”

Book of Abstracts

June 20-21, 2019
Belgrade, Serbia



Scientific committee

Tanja Cirkovic Velickovic (President)

Paola Roncada

Andrea Urbani

Irena Vovk

Bruno De Meulenaer

Karin Hoffmann-Sommergruber

Marianne van Hage

Tatjana Parac-Vogt

Andreja Rajkovic

Michelle Epstein

Danijela Apostolovic

Organizing Committee

Sanja Grguric-Sipka (Chair)

Mirjana Macvanin (Co-Chair)

Dusanka Milojkovic-Opsenica

Maja Mladenovic

Teodora Djukic

Tatjana Bozic

Mirjana Radomirovic



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 810752

P4

Physicochemical characterization of soluble proteins of whole camel milk powders produced by spray drying treatment at high temperatures

*Marija Peruško*¹, Ana Simović², Nikola Stevanović³, Katarina Smiljanić⁴, Mirjana Radomirović⁴, Dragana Stanić-Vučinić⁴, Sami Ghnimi^{5,6,7} and Tanja Ćirković Veličković^{4,5,6,8}

¹Faculty of Chemistry - Innovation Center d.o.o., Belgrade, Serbia; ²Department of Biochemistry, University of Belgrade - Faculty of Chemistry, Belgrade, Serbia; ³Department of Analytical Chemistry, University of Belgrade - Faculty of Chemistry, Belgrade, Serbia; ⁴Center of Excellence for Molecular Food Sciences & Department of Biochemistry, University of Belgrade - Faculty of Chemistry, Belgrade, Serbia; ⁵Ghent University Global Campus, Yeonsu-Gu, Incheon, South Korea; ⁶Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium; ⁷Department of Food Science, United Arab Emirates University, UAE; ⁸Serbian Academy of Sciences and Arts, Belgrade, Serbia

Objective. Camel milk is highly nutritious food with numerous health benefits proposed. Demand for camel milk has increased worldwide. Production of camel milk powders facilitate its transport, prolonge shelf-life, and also offer an attractive additive for various food products. In this study we characterized proteins of soluble fraction of freeze/spray dried camel milk powders.

Material and Methods. Whole camel milk powders were prepared by spray drying treatment at six different inlet temperatures (190°C - 250°C) or by freeze drying. The soluble protein fractions upon the treatments were analysed by combination of electrophoretic and spectroscopic techniques. Functional properties, such as antioxidant activity and protein solubility were assessed.

Results. SDS-PAGE revealed non-uniform increase in Mw of major protein bands, while native electrophoresis revealed non-uniform decrease in pI values with increased inlet temperature of spray drying. That indicated attachment of lactose moieties to NH₂-group of proteins via non-enzymatic Maillard reaction. Spectrophotometric analysis showed formation of intermediate Maillard reaction products (increased absorbance at 294 nm) and no detectable late Maillard reaction products formation. Far-UV circular dichroism spectra showed no differences in secondary structures between freeze and spray dried samples. Antioxidant activity and protein solubility were increased with increase in inlet temperature.

Conclusions. Our results showed that spray drying treatment promoted non-enzymatic glycation of camel milk proteins. Glycation of food proteins affects their techno-functional properties, shelf-life and nutritional value. Thus, optimization of spray drying parameters is essential for production of high quality camel milk powders.

Acknowledgements: This research work was funded the Ministry of Education and Science of the Republic of Serbia, GA No. OI172024, Ghent University Global Campus, Belgian Special Research Fund BOF StG No. 01N01718, Serbian Academy of Sciences and Arts Project F-26. The project leading to this application has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 810752.

Београд, 31. октобар, 2019.

Проф. др Тања Ћирковић Величковић
Хемијски факултет
Универзитет у Београду

Поштована др Ћирковић Величковић,

Разматран је захтев за евалуацију категорије научног скупа, који сте доставили 12. јула 2019., писаним путем Матичном одбору за хемију

1st FoodEnTwin Workshop “Food and Environmental -Omics”, одржаног 20-21. јуна 2019, у Београду, у организацији Хемијског факултета, Универзитета у Београду.

Одбор је донео одлуку да према критеријумима из важећег Правилника о поступку и начину вредновања и квантитативном исказивању научноистраживачких резултата истраживача, научни скуп је

међународног карактера.

С поштовањем,



Проф. др Живослав Тешић
Председник Матичног научног одбора за хемију