

Supplementary data for article:

Malenov, D. P.; Ninkovic, D. B.; Zaric, S. D. Stacking of Metal Chelates with Benzene:
Can Dispersion-Corrected DFT Be Used to Calculate Organic-Inorganic Stacking?
ChemPhysChem **2015**, *16* (4), 761–768. <https://doi.org/10.1002/cphc.201402589>

CHEMPHYSICHEM

Supporting Information

Stacking of Metal Chelates with Benzene: Can Dispersion-Corrected DFT Be Used to Calculate Organic–Inorganic Stacking?

Dušan P. Malenov,^[c] Dragan B. Ninković,^[c] and Snežana D. Zarić*^[a, b]

cphc_201402589_sm_miscellaneous_information.pdf

SUPPORTING INFORMATION




Figure S1. Performance of dispersion corrected GGA, H-GGA and M-GGA functionals for stacking interactions of metal-chelates with benzene, with def2-TZVP and aug-cc-pVDZ basis sets, and without BSSE correction; black lines within every system denote the benchmark CCSD(T) energy




Figure S2. Performance of dispersion corrected HM-GGA, LC-GGA and DH-GGA functionals and MP2 method for stacking interactions of metal-chelates with benzene, with def2-TZVP and aug-cc-pVDZ basis sets, and without BSSE correction; black lines within every system denote the benchmark CCSD(T) energy