

Supplementary data for the article:

Filipović, N. R.; Bjelogrić, S.; Todorović, T. R.; Blagojević, V. A.; Muller, C. D.; Marinković, A.; Vujčić, M.; Janović, B.; Malešević, A. S.; Begović, N.; et al. Ni(II) Complex with Bishydrazone Ligand: Synthesis, Characterization, DNA Binding Studies and pro-Apoptotic and pro-Differentiation Induction in Human Cancerous Cell Lines. *RSC Adv.* **2016**, *6* (110), 108726–108740. <https://doi.org/10.1039/c6ra24604d>

Electronic Supplementary Information

**Ni(II) complex with bishydrazone ligand: synthesis, characterization, DNA binding studies
and pro-apoptotic and pro-differentiation induction in human cancerous cell lines**

Nenad R. Filipović^{a,*}, Snežana Bjelogrić^b, Tamara R. Todorović^c, Vladimir A. Blagojević^d,
Christian D. Muller^e, Aleksandar Marinković^f, Miroslava Vujčić^g, Barbara Janović^g, Aleksandar
S. Malešević^c, Nebojša Begović^h, Milan Senčanskiⁱ, Dragica M. Minić^j

^a*Faculty of Agriculture, University of Belgrade, Nemanjina 6, 11000 Belgrade, Serbia;*

^b*National Cancer Research Center of Serbia, Pasterova 14, 11000 Belgrade, Serbia.*

^c*Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, 11000 Belgrade, Serbia*

^d*Institute of Technical Sciences SASA, Knez Mihailova, 35/IV, 11000 Belgrade, Serbia*

^e*Institut Pluridisciplinaire Hubert Curien, UMR 7178 CNRS Université de Strasbourg, 67401
Illkirch, France*

^f*Faculty of Technology and Metallurgy, University of Belgrade, Karnegijeva 4, 11000 Belgrade,
Serbia*

^g*Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoševa 12,
11000 Belgrade, Serbia*

^h*Institute of General and Physical Chemistry, Studentski trg 12, 11000 Belgrade, Serbia*

ⁱ*Center for Multidisciplinary Research, Institute of Nuclear Sciences "Vinča", University of
Belgrade, 11000 Belgrade, Serbia*

^j*Faculty of Physical Chemistry, University of Belgrade, Studentski trg 12-16, 11000 Belgrade,
Serbia*

***Corresponding author**

Nenad R. Filipović

Faculty of Agriculture

University of Belgrade

Nemanjina 6, 11000 Belgrade, Serbia

Email: nenadf.chem@gmail.com

Content

Figure S1. Mass spectrum of 1 in acetonitrile	3
Figure S2. ^1H NMR spectrum of 1 in CDCl_3	4
Figure S3. ^{13}C NMR spectrum of 1 in CDCl_3	5
Figure S4. COSY spectrum of 1 in CDCl_3	6
Figure S5. NOESY spectrum of 1 in CDCl_3	7
Figure S6. ROESY spectrum of 1 in CDCl_3	8
Figure S7. ^1H - ^{13}C HSQC NMR spectrum of 1 in CDCl_3	9
Figure S8. ^1H - ^{13}C HMBC NMR spectrum of 1 in CDCl_3	10
Table S1. Experimental and calculated ^1H NMR chemical shifts of 1	11
Table S2. Experimental and calculated ^{13}C NMR chemical shifts of 1	12
Figure S9. Cyclic voltammograms of 1 in anhydrous DMSO	13
Figure S10. Concentration-response curve and ED_{50} value for 1 on THP-1 cells treated during 24 h incubation	13




Figure S1. Mass spectrum of **1** in acetonitrile.




Figure S2. ^1H NMR spectrum of **1** in CDCl_3 .




Figure S3. ^{13}C NMR spectrum of **1** in CDCl_3 .




Figure S4. COSY spectrum of **1** in CDCl₃.




Figure S5. NOESY spectrum of **1** in CDCl₃.




Figure S6. ROESY spectrum of **1** in CDCl₃.




Figure S7. ^1H - ^{13}C HSQC NMR spectrum of **1** in CDCl_3 .




Figure S8. ^1H - ^{13}C HMBC NMR spectrum of **1** in CDCl_3 .

Table S1. Experimental and calculated ^1H NMR chemical shifts of **1**.

Assignment	Experimental chemical shift δ (ppm)	Calculated chemical shift δ (ppm)
H-C3	8.44	8.22577
H-C4	7.83	7.86877
H-C2	7.97	7.68497
H-C7	2.29	2.38157
H-C14	2.43	2.57017
H-C9	3.44	3.13757
H-C16	3.72	4.10837
H-C11	4.25	4.00027
H-C18	4.17	4.14927
H-C12	1.33	1.31587
H-C19	1.29	1.32397
H ₂ O	2.11	2.31637
		9.68517

Table S2. Experimental and calculated ^{13}C NMR chemical shifts of **1**

Assignment	Experimental chemical shift δ (ppm)	Calculated chemical shift δ (ppm)
C3	141.60	136.0176
C4	123.19	118.5201
C2	124.64	119.9311
C5	156.28	150.5638
C1	137.80	139.1615
C6	153.08	158.8117
C13	131.57	138.3357
C7	12.06	16.3772
C14	20.98	22.2051
C8	182.10	166.0884
C15	178.04	175.0396
C9	37.96	45.1723
C16	45.48	50.8007
C10	168.66	170.9611
C17	169.76	165.6791
C11	61.29	64.2114
C18	60.58	63.5428
C12	14.18	14.3748
C19	14.11	14.4742




Figure S9. Cyclic voltammograms of **1** in anhydrous DMSO containing 0.10 M $[n\text{-Bu}_4\text{N}][\text{PF}_6]$ at a scan rate of 100 mVs^{-1} using a glassy carbon working electrode in the potential range from -2.5 to 0.0 V (A) and from 0.0 to $+1.5$ V (B).




Figure S10. Concentration-response curve and ED_{50} value for **1** on THP-1 cells treated during 24 h incubation. Results are presented as percentage of apoptotic events determined by means of Annexin V/PI double staining in two independent experiments (circles and crosses) with asymmetric five-parameter sigmoidal curve computed for both replicated in GraphPad Prism software.




Figure S11. Comparison of experimental IR spectrum and calculated IR spectrum for the structure with coordination of the carbonylic oxygen of the hydrazonic arm.




Figure S12. UV-VIS spectra of **1** in DMSO over 24 hour period.




Figure 13. DSC curves in air with subtracted baselines at different heating rates (A), with enhanced region of the first degradation peak (B).




Figure S14. DSC recorded in nitrogen at lower (A) and higher (B) heating rates.