

Supplementary data for the article:

Pantić, D. N.; Arancrossed D Sigelović, S.; Radulović, S.; Roller, A.; Arion, V. B.; Grgurić-Šipka, S. Synthesis, Characterisation and Cytotoxic Activity of Organoruthenium(II)-Halido Complexes with 1H-Benzimidazole-2-Carboxylic Acid. *Journal of Organometallic Chemistry* **2016**, *819*, 61–68.
<https://doi.org/10.1016/j.jorganchem.2016.06.024>

Supplementary Materials for:

Synthesis, characterisation and cytotoxic activity of organoruthenium(II)-halido complexes with 1*H*-benzimidazole-2-carboxylic acid

Darko N. Pantić^a, Sandra Arandelović^b, Siniša Radulović^b, Alexander Roller^c, Vladimir B. Arion^c, Sanja Grgurić-Šipka^{a,*}

^a Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, 11000 Belgrade, Serbia

^b Institute for Oncology and Radiology of Serbia, Pasterova 14, 11000 Belgrade, Serbia

^c Institute of Inorganic Chemistry, University of Vienna, Währinger Str. 42, 1090 Vienna, Austria

* Corresponding author. *E-mail*: sanjag@chem.bg.ac.rs.

Contents:

1. Fig. S1–S3 FTIR spectra of C1–C3	S2
2. Fig. S4–S6 ¹ H NMR spectra of C1–C3	S3
3. Fig. S7–S9 ¹³ C NMR spectra of C1–C3	S5
4. Fig. S10–S12 ESI-mass spectra of C1–C3	S6
5. Fig. S13 and S14 UV–vis spectra of C1	S8
6. Fig. S14 UV–vis spectra of C3	S8
7. Table S1 Rate Constants for the aquation at 298 K for complex C1 and C3	S9




Fig. S1. FTIR spectra for C1.




Fig. S2. FTIR spectra for C2.




Fig. S3. FTIR spectra for C3.




Fig. S4. ¹H NMR spectra for C1.




Fig. S5. ^1H NMR spectra for C2.




Fig. S6. ^1H NMR spectra for C3.




Fig. S7. ^{13}C NMR spectra for C1.




Fig. S8. ^{13}C NMR spectra for C2.




Fig. S9. ^{13}C NMR spectra for C3.




Fig. S10. ESI-MS spectra for C1.




Fig. S11. ESI-MS spectra for C2.




Fig. S12. ESI-MS spectra for C3.




Fig. S13. UV-vis spectra of complex **C1** in water-PEG400 mixture (85 : 15) soon after dissolution (black line), after 0.5 h (red line), after 1 h (green line) and after 24 h (blue line) incubation at 298 K.




Fig. S14. UV-vis spectra of complex **C3** in water-PEG400 mixture (85 : 15) soon after dissolution (black line), after 0.5 h (red line), after 1 h (green line) and after 24 h (blue line) incubation at 298 K.

Table S1 Rate Constants for the aquation at 298 K in water-PEG400 mixture (85 : 15) for **C1** and **C3**.

Compound	k_{obs} (10^{-5} , s^{-1})	$t_{1/2}$ (s)
C1	60.4225 ± 2.47049	1102
C3	51.1243 ± 6.744	1198